



Gut Morphology and Internal Organs of Broiler Birds Fed Graded Levels of Bio-Detheobrominized Cocoa Bean Shell (CBS) Based Diets

G. O. Adeyemo^{1*}, A. O. Ajayi¹, O. G. Longe¹ and O. Olubamiwa²

¹Department of Animal Science, Faculty of Agriculture, University of Ibadan, Nigeria.

²Department of Animal Science, College Agricultural Sciences, Landmark University, Omu-Aran, Kwara State, Nigeria.

Authors' contributions

This work was carried out in collaboration between all authors. Author OGL designed the concept of the study. Author OO wrote the protocol and reviewed the experimental design. Author GOA supervised the feeding trials, wrote the first draft of the manuscript and all drafts of the manuscript. Author AOA carried out the feeding trials, managed the analyses of the study and was involved in contributing to the final drafts of manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJEA/2015/10582

Editor(s):

(1) Ismail Seven, Department of Plant and Animal Production, Vocational School of Sivrice, University of Firat, Turkey.

Reviewers:

(1) Anonymous, Usmanu Danfodiyo University, Sokoto.

(2) Ndubueze Amaechi, Department of Veterinary Microbiology and Parasitology, Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria.

Complete Peer review History: <http://www.sciencedomain.org/review-history.php?iid=665&id=2&aid=6316>

Original Research Article

Received 1st April 2014
Accepted 5th September 2014
Published 4th October 2014

ABSTRACT

The effects of feeding graded levels of bio-detheobrominized cocoa bean shell (CBS) based diets were studied in broilers. Detheobromination by fungi fermentation (bio-detheobromination) reduces the methylxanthine contents cocoa by-products when fermented with *Aspergillus niger*. 196 day old broiler chicks were used for the experiment in an eight week feeding trial. There were seven dietary treatments containing graded levels of CBS from 0 to 30% at the expense of maize and soyabean meal. The control group had no CBS in its diet. Parameters measured include feed intake, body weight gain, feed conversion ratio, carcass characteristics and gut morphology.

The results showed that total feed intake, average body weight gain, feed conversion ratio, weight of internal organs and gut morphology were significantly ($p < 0.05$) different across dietary treatments while the carcass characteristics did not show any significant ($p > 0.05$) difference. The result of the study revealed that there was a significant reduction in the weight gain and feed efficiency of the

undisturbed to allow the growth of the mycelia through the substrates. The mycelia were grown for eight weeks on the substrate after which the substrate were dried and milled to homogeneity. One hundred grams of sterile CBS powder (Substrate) were inoculated with 1 gram *Aspergillus niger* spore. Substrate inoculated with the *A. niger* spores were kept in a room temperature condition for seven days after which diets containing graded levels of fermented CBS were formulated and compounded.

2.3 Experimental Design and Management

One hundred and ninety six (196) Day old Ross broiler chicks were randomly allotted to seven (7) dietary treatments in a completely randomized design with four replicates and seven birds per replicate. The diets shown in Tables 2 and 3 contained graded levels of CBS at 0, 5, 10, 15, 20, 25 and 30% respectively. Diet one had no CBS and served as the control. The birds were reared on deep litter. The usual conventional brooding practices and poultry management were observed during the experimental period. Routine vaccination schedule were properly observed and the health of the birds were well taken care of.

2.4 Data Collection

One bird per replicate in each of the seven treatment diets (Tables 2 and 3) were randomly selected at eight weeks old, slaughtered, dressed and weighed to determine warm dressed weight. The carcass was eviscerated and the internal organs separated and weighed (expressed as percentage of live-weight). Duodenum and ileal sections of each sacrificed bird were cut to determine and compare their villus height, crypt depth & villus width.

2.5 Gut Morphological Studies

The entire intestine of each sacrificed bird was removed and 5cm intestinal sections from the duodenum (after the duodenal loop) and ileum (5cm before the cecal tonsils) were collected. Sections were cleared and embedded in paraffin where they were initially fixed in 10% phosphate-buffered formalin for 24 hours, transferred to 50% ethanol solution for 15 minutes, and immersed in 70% ethanol. Formalin-fixed sample were embedded in paraffin and 5-um section were sliced and stained with heamatoxylin. The

samples were analyzed under a standard light microscope with special emphasis on gut architecture as described by Brady et al. [10]. The villous height, villous width and crypt depth were determined using computer aided light microscope image analysis as described by [11]. Sections from one bird per replicate in each of the seven treatments were placed on each slide. A total of 5 digital images were taken from each tissue using a microscope with digital camera. From each slide, 5 areas were chosen, and 10 villi from each of these areas were evaluated for villus height and width. In addition, 10 crypts were evaluated for crypt depth at each of these areas.

2.6 Proximate Composition of CBS and Diets

Proximate composition of the fermented CBS and experimental diets shown in Table 1 (crude protein, crude fiber, ether extract, ash) were determined using the procedure of AOAC [12]. The anti-nutritional factor (theobromine) was determined using gravimetric method by Holmes [15].

2.7 Statistical Analysis

All data obtained were analyzed using analysis of variance of the completely randomized design [13] and least significant difference were used to separate the treatment means.

3. RESULTS AND DISCUSSION

3.1 Internal Organ Weight (% of live weight) of Birds Fed CBS-Based Diets

Results of the internal organ weights of birds fed CBS- based diets are presented in Table 4. It was observed that there was no significant difference ($P>0.05$) in the values obtained for liver and heart. However, lower weight was observed at 5%, while the higher weight was observed at 25%. There were however significant differences ($P<0.05$) for the values obtained for gizzard, spleen, pancreas and abdominal fat pad weights. Gizzard weights revealed that the lowest was observed at 10% and the highest weight at 30%, and this same trend was observed for pancreas weights also, this could be a reflection of the high fiber content of the feed, which leads to increased organ weights as a result of work which is required in the digestion. Abdominal fat pad weights were

lower at 20% inclusion levels and higher on the control diet. This could be as a result of the nutrient diluting effect of fiber in the diets of monogastrics.

3.2 Gut Morphology of Broiler Birds Fed Graded Levels of CBS Based Diets

Presented in Table 5 are the results of gut morphology analysis of broiler birds fed graded levels of CBS-based diets. At both the duodenum and the ileum, significant differences were observed for villus height, crypt depth and villus: crypt ratio for birds fed the graded levels of CBS in their diets, with the results indicating that the villous height in duodenum was greater than those in the ileum for broiler birds fed varying levels of CBS-based diets. This observation is in agreement with the findings of Gholamhazan et al. [14] who observed that villi height in duodenum was greater than those in the jejunum and ileum when broiler chickens were fed diets varying in the levels of lysine. The finding above is consistent to the major role of duodenum in nutrient absorption. The villous height depends upon two factors, firstly on cellular multiplication and differentiation from the crypts and secondly on the migration of mature cells along the villous axis, accompanied by the cellular shedding from the apex. The regulation of these factors is

widely considered to be the way in which the small intestine adapts to nutritional demands [15]. The duodenum villous height and crypt depth of broiler birds fed graded-levels of CBS-based diets increased significantly ($P < 0.05$) in the CBS-diets compared to the control. However the increased in the villous height across the CBS diets did not result in increased weight gain and feed utilization. Increased villi height is expected to result in improved performance as a result of improved nutrient absorption. The increase in villi height observed in the CBS diets might indicate that the birds fed CBS diets up to 30% CBS dietary level might have had greater nutrient absorption and utilization because increase in villi height implies increased surface area for nutrient utilization [16].

Table 1. Chemical composition of cocoa bean shell

Component	Values
Dry matter (%)	84.52
Crude Protein (%)	14.98
Ether Extract (%)	2.39
Crude Fiber (%)	7.67
Ash (%)	11.66
Metabolizable Energy (kcal/kg)	2,800
Theobromine (mg/g)	50

Table 2. Composition of CBS based diets fed to broiler birds at starter phase (Kg/100kg) diets

Ingredients	0%CBS	5%CBS	10%CBS	15%CBS	20%CBS	25%CBS	30%CBS
CBS	0.00	5.00	10.00	15.00	20.00	25.00	30.00
Maize	53.00	50.00	46.00	43.00	40.00	37.00	34.00
Soya Bean Meal	38.00	36.00	35.00	33.00	31.00	29.00	26.00
Fish Meal	3.90	3.90	3.90	3.90	3.90	3.90	3.90
Bone Meal	2.00	2.00	2.00	2.00	2.00	2.00	2.00
DCP	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Limestone	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Salt	0.25	0.25	0.25	0.25	0.25	0.25	0.25
B-Premix	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Lysine	0.15	0.15	0.15	0.15	0.15	0.15	0.15
Methionine	0.20	0.20	0.20	0.20	0.20	0.20	0.20
TOTAL	100.00	100.00	100.00	100.00	100.00	100.00	100.00
Proximate Analysis							
M.E(kcal/kg)	3,185.38	3,156.42	3,126.14	3,097.18	3,068.22	3,039.26	3,010.33
Crude Protein (%)	24.46	24.04	23.96	23.54	23.11	22.69	22.26
Ether Extract (%)	3.47	3.28	3.09	2.89	2.71	2.52	2.33
Crude Fiber (%)	3.55	3.76	3.80	4.03	4.84	4.95	5.21
Calcium (%)	1.20	1.55	1.79	2.04	2.29	2.53	2.46
Phosphorus (%)	0.61	0.54	0.68	0.82	0.96	1.09	1.23

M.E= Metabolisable Energy (kcal/kg)

Table 3. Composition of CBS based diets fed to broiler birds at finisher phase (kg/100kg) diets

Ingredients	0%CBS	5%CBS	10%CBS	15%CBS	20%CBS	25%CBS	30%CBS
Cocoa Bean Shell	0.00	5.00	10.00	15.00	20.00	25.00	30.00
Maize	63.00	60.00	57.00	54.00	51.00	48.00	45.00
Soya Bean Meal	31.00	29.00	27.00	25.00	23.00	21.00	19.00
Fish Meal	0.70	0.70	0.70	0.70	0.70	0.70	0.70
Molasses	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Bone Meal	1.50	1.50	1.50	1.50	1.50	1.50	1.50
DCP	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Limestone	1.0	1.0	1.00	1.00	1.00	1.00	1.00
Salt	0.25	0.25	0.25	0.25	0.25	0.25	0.25
B-Premix	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Lysine	0.15	0.15	0.15	0.15	0.15	0.15	0.15
Methionine	0.15	0.15	0.15	0.15	0.15	0.15	0.15
TOTAL *	100.00	100.00	100.00	100.00	100.00	100.00	100.00
Proximate Analysis							
M.E(kcal/kg)	3,224.93	3,195.97	3,167.01	3,138.05	3,109.09	3,080.13	3,051.71
Crude Protein(%)	20.31	19.89	19.46	19.04	18.61	18.91	18.76
Ether Extract(%)	3.62	3.42	3.24	3.05	2.86	2.67	2.48
Crude Fiber(%)	3.29	3.40	3.80	4.55	4.90	5.26	5.50
Calcium(%)	0.81	1.06	1.31	1.55	1.79	2.02	2.29
Phosphorus(%)	0.45	0.52	0.65	0.79	0.92	1.06	1.98

Table 4. Weights of internal organs (%) of broilers fed graded-levels of CBS based diets

Parameters	0%CBS	5%CBS	10%CBS	15%CBS	20%CBS	25%CBS	30%CBS
Liver	1.793	1.673	1.983	2.055	2.195	2.25	2.058
Gizzard	3.12 ^b	2.995 ^d	2.915 ^b	3.258 ^{ab}	3.253 ^{ab}	3.558 ^{ab}	3.728 ^a
Spleen	0.022 ^{bc}	0.034 ^{bc}	0.067 ^{ab}	0.094 ^a	0.179 ^c	0.088 ^{ab}	0.051 ^{abc}
Heart	0.4363	0.4875	0.4675	0.4675	0.565	0.465	0.565
Pancreas	0.162 ^c	0.258 ^{ab}	0.158 ^c	0.220 ^b	0.250 ^{ab}	0.275 ^{ab}	0.283 ^a
Abdominal fat	1.24 ^a	0.55 ^{bc}	0.99 ^{ab}	0.86 ^{ab}	0.23 ^c	0.59 ^{bc}	0.99 ^{ab}

Table 5. Gut morphology of birds fed CBS based diets

Parameters	0%CBS	5%CBS	10%CBS	15%CBS	20%CBS	25%CBS	30%CBS	SEM
Villus height								
Duodenum	1589 ^{bc}	1681 ^b	1740 ^b	1176 ^c	2074 ^a	1210 ^c	2402 ^a	8.10
Ileum	974.9 ^c	1806 ^a	1180 ^{bc}	1537 ^{ab}	1767 ^a	1392 ^b	135.1 ^d	3.7
Villus width								
Duodenum	165.2 ^c	212.5 ^b	219.2 ^b	192.8 ^d	148.1 ^c	153.1 ^c	253.1 ^a	5.193
Ileum	208.7 ^{ab}	228.5 ^b	191.3 ^b	225.7 ^{ab}	258.2 ^a	208.1 ^{ab}	233.5 ^{ab}	19.58
Crypt depth								
Duodenum	66.73 ^{cd}	149.7 ^b	45.49 ^d	98.06 ^c	189.6 ^b	301.8 ^a	152.4 ^b	3.94
Ileum	93.79 ^c	235.8 ^a	119.9 ^c	142.6 ^d	214.5 ^a	134.5 ^b	136.8 ^b	10.6

^{abc}Means with the same super script are not significantly different ($P>0.05$)

4. CONCLUSION

The findings in this study suggest that although the gut morphology of the birds fed CBS diets improved compared to the control diet, to have enhanced nutrient absorption, essential nutrients such as digestive protein and energy were locked up or bonded by anti-nutritional factors such as theobromine and tannins present in the CBS diets thereby impairing the essential nutrient's bio-availability thereby impeding the

growth of the birds fed diets beyond the 10% inclusion level. The results of this study showed that increasing the dietary level of cocoa bean shell in broiler chicken's diets improved their villous and crypt dimensions positively.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Aina AA. Potential of cocoa by-product as livestock feeds. Proceeding of National Workshop on Alternative Formulation of Livestock Feed held at ARMTI, Ilorin. 1998;21-25.
2. Carolien M. Acid binding capacity in feedstuffs. Feed Int. October. 2001;21-25.
3. Odunsi AA, Longe OG. The nutritive value of hot water or cocoa pod ash solution treated cocoa bean cake for broiler chicks. Brit. Poult. Sci. 1998;39:519-525.
4. Olubamiwa O, Soetan OA, Olamijulo OA, Hamzat RA, Longe OG. Utilization of variously treated cocoa bean shells in layer mash. Proc., 27th Annual Conf., Nig. Soc. of Animal Production (NSAP) March 17-21, 2002. Fed. University of Technology, Akure, Nigeria. 2002;267-269.
5. Olupona JA, Abodunwa JA, Fayoyin FK. Response of laying hens to graded levels of cocoa bean shells. In: Proceedings of the 28th Ann. Conf., Nig. Soc. Anim. Prod. (NSAP). 2003;28:247-249.
6. FAO. Animal feed resources information system. Tropical Feeds. 2002;76-82.
7. Olubamiwa O, Hamzat RA. Effect of processing techniques on the utilization of cocoa bean shell by laying hens. Proc. Malaysian International Cocoa Conference, July 18-19, 2005, Kuala Lumpur, Malaysia. 2005;118-119.
8. Alemawor F, Dzogbefia VP, Oldham JH, Oddoye OK. Effect of fermentation on the composition of cocoa pod husk: Influence on time and Mn²⁺ supplementation on the fermentation process. African Journal of Biotechnology. 2009;8(9):1950-1958.
9. Adamafio NA, Ayombil F, Tano-Debrah K. Microbial de-theobromination of cocoa (*Theobroma cacao*) pod husk. Asian J. Biochem 2011;6:200-207.
10. Brady J, Hernandez-Doria JD, Rodriguez-Lecompte JC. Toxinotyping of necrotic enteritis-producing and commensal isolates of *Clostridium perfringens* from chicken fed organic diets. Avian pathol. 2010;39:475-481
11. Bird TA, Kyriakis JM, Gayle M, Milne A, Virca D. Interleukin-1 activates P54 nitrogen-activated protein (MAP) Kinase/ Stress activated protein Kinase by a pathway that is independent of P 21ras, Raf-1 and MAP Kinase. J. Biol Chem. 269 J.D, Benneth C, Gudiquez- Lemcompte JC 2010. Toxinotyping of necrotic enteritis-producing and commensal isolates of *Clostridium perfringens* from Chickens fed organic diets. Avian Pathol. 1994;39:475-481.
12. Association of Official Analytical Chemists AOAC. Methods of Analysis, 15th edition, Arlington, VA. 2002;4:54.
13. SAS. Statistical Analysis System Institutes. User's guide SAS Institute Inc. version 10 edition. Cary, North Carolina; 2010.
14. Gholamhasan V, Teshfam M, Bahadoran S, Farazyan H, Hossein S. Effects of different levels of lysine on small intestinal villous morphology in starter diets of broiler chickens. Global Veterinaria. 2011;7(6):523-526.
15. Holmes KE. Determination of theobromine in cocoa products. Analyst. 1950;75:457-461.
16. Ofusori DA, Caxton-Martins EA. A Comparative histomorphometric study of the stomach of Rat (*Rattus norvegicus*), Bat (*Eidolon helvum*) and Pangolin (*Manis tricuspis*) in relation to diet. Int. J. Morphol. 2008;26(3):669-674.

© 2015 Adeyemo et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history.php?iid=665&id=2&aid=6316>